

PROPOSITION 4. *If R is semiprime and $x \in R$, then x has LS and RS simultaneously; if some x in R has both LB and RB, then R is unital.*

Thus for a semiprime R there are five possible combinations: \emptyset , $\{LB\}$, $\{RB\}$, $\{LS, RS\}$, Δ .

Full details will appear in [3].

REFERENCES

- 1 B. A. Barnes, The Fredholm elements of a ring, *Canad. J. Math.* 21:84–95 (1969).
- 2 B. A. Barnes, G. J. Murphy, M. R. F. Smyth, and T. T. West, *Riesz and Fredholm Theory in Banach Algebras*, Pitman Research Notes in Mathematics No. 67, 1982.
- 3 M. R. F. Smyth and T. T. West, Barnes and support idempotents associated with ring elements, *Proc. Roy. Irish Acad. Sect. A*, to appear.

Products of Skew-Symmetric Matrices

Thomas J. Laffey
Department of Mathematics
University College Dublin
Belfield
Dublin 4, Ireland

This note summarizes some recent results obtained jointly by R. Gow and the author. Detailed proofs will appear elsewhere.

Throughout this note, F denotes an algebraically closed field of characteristic not equal to 2, and $A \in M_n(F)$ is a nonsingular $n \times n$ matrix.

We consider the following questions:

- (1) Can A be expressed as a product of a finite number k of skew-symmetric matrices in $M_n(F)$?
- (2) If so, what is the smallest number k for which such a representation is possible?

Note that since A is nonsingular, a necessary condition is that n is even. Also, the product of two skew-symmetric 2×2 matrices is scalar, so a 2×2 matrix A satisfies (1) if and only if it is either skew-symmetric or scalar.

We now state our main results.

THEOREM 1 [2]. *A is a product of two skew-symmetric matrices if and only if A is similar to a matrix of the form*

$$\begin{bmatrix} B & 0 \\ 0 & B \end{bmatrix}.$$

THEOREM 2. *Suppose $n \equiv 0 \pmod{4}$. Then A can be expressed as a product of five skew-symmetric matrices. There is a nonsingular matrix A which cannot be expressed as a product of fewer than five skew-symmetric matrices.*

THEOREM 3. *Suppose $n > 2$ and $n \equiv 2 \pmod{4}$. Then A can be expressed as a product of seven skew-symmetric matrices.*

REMARK. We do not know whether "seven" can be replaced by "five" in Theorem 3.

We now sketch the proof of Theorem 2. By a result of Gow [1], $A = SJ$, where $S \in M_n(F)$ is symmetric and J is an involution, i.e., $J^2 = I$. A careful analysis of the proof shows that J may be chosen to have $n/2$ eigenvalues equal to $+1$ and $n/2$ eigenvalues equal to -1 . Since $n \equiv 0 \pmod{4}$, it thus follows that J is similar to the matrix of the form

$$\begin{bmatrix} B & 0 \\ 0 & B \end{bmatrix}, \quad \text{where } B = \begin{bmatrix} I_{n/4} & 0 \\ 0 & -I_{n/4} \end{bmatrix},$$

and hence, by Theorem 1, J is a product of two skew-symmetric matrices. Next, since F is algebraically closed, there is a nonsingular matrix P in $M_n(F)$ such that

$$P'SP = I.$$

Let K be a nonsingular skew-symmetric matrix in $M_{n/2}(F)$. Then

$$\begin{bmatrix} K^{-1} & 0 \\ 0 & K^{-1} \end{bmatrix}$$

is a product of two skew-symmetric matrices by Theorem 1, and hence $I = S_1 S_2 S_3$ is a product of three skew-symmetric matrices. Hence

$$S = (P'^{-1} S_1 P^{-1})(P S_2 P')(P'^{-1} S_3 P^{-1})$$

is a product of three skew-symmetric matrices, and thus A is a product of five skew-symmetric matrices.

We next show that there is a nonsingular matrix A in $M_n(F)$ (n even) which cannot be expressed as a product of four skew-symmetric matrices. Suppose $A = P_1 P_2 P_3 P_4$ where P_1, P_2, P_3, P_4 are skew-symmetric. By Theorem 1, $A = XY$ where X, Y have each of their elementary divisors occurring with even multiplicity. Taking A to be a diagonal matrix $\text{diag}(1, 1, \dots, 1, a)$ where $a \neq 0, 1$, we use the Sá-Thompson interlacing theorem for invariant factors [3, 4] to get a contradiction. This shows that there is an A which is not a product of four skew-symmetric matrices. Hence there is a nonsingular A_0 which cannot be written as a product of three skew-symmetrics (for example, take $A_0 = AK^{-1}$ where K is a skew-symmetric and A cannot be written as a product of four skew-symmetric matrices). It then follows that there is a nonsingular A_1 which cannot be expressed as a product of fewer than five skew-symmetric matrices.

The proof of Theorem 3 is similar in style to that of Theorem 2, but more difficult because J does not satisfy Theorem 1. The key step is to show that if $K \in M_n(F)$, $n > 2$, $n \equiv 2 \pmod{4}$, is nonsingular and skew-symmetric, then K can be expressed as a product of four skew-symmetrics. This is done by examination of the Jordan form of K ; an inductive argument reduces the problem to two cases:

- (1) K similar to $J_{n/2}(\lambda) \oplus J_{n/2}(-\lambda)$,
- (2) K similar to

$$J_{(n-2)/2}(\lambda) \oplus J_{(n-2)/2}(-\lambda) \oplus \begin{bmatrix} 0 & a \\ -a & 0 \end{bmatrix}$$

[where $J_r(\lambda)$ denotes the $r \times r$ Jordan block with eigenvalue λ], and these are dealt with by direct computation.

We have extended some of the results to more general fields F (assuming $\det A$ is a square in F)—the details will appear elsewhere.

REFERENCES

- 1 R. Gow, The equivalence of an invertible matrix to its transpose, *Linear and Multilinear Algebra* 8:329–336 (1980).
- 2 R. Gow and T. J. Laffey, Pairs of alternating forms and products of two skew symmetric matrices, *Linear Algebra Appl.* 63:119–132 (1984).
- 3 E. Marques de Sá, Imbedding conditions for λ -matrices, *Linear Algebra Appl.* 24:33–50 (1979).
- 4 R. C. Thompson, Interlacing inequalities for invariant factors, *Linear Algebra Appl.* 24:1–31 (1979).